

# Flora of Maine

List of plants observed by J. S. Collins, Malden, Mass.

C = Cape Royer, Maine, July 10-25, 1892

E = Eagle Island " " 15-21 "

S. = Great Spruce Head " " 17. "

Numbers from Portland Catalogue, 2d. Edition, 1892

#12	C	291 C
16	C E	292 C
28	C E S	293 C E S
31	C	299 C
33	C	300 C E S
37	E	306 E
47	E	312 C
52	E	313 C S
78	E	325 C
81	E	333 C E S
82	C	352 C E
83	C E S	354 C
88	E	359 E
96a	C E	360 C E S
128	C	363 C E S
130	C E S	370 E
133	C	371 E
138	C E S	374 C E
140	C E S	379 C E S
143	C E	380 C E
145	C E	385 C E S
152	C E	387 C
155	C S	389 C
157	E	392 C S
159	S. near old house	400 S
170	(E. S. also at E, a form with perfectly white flowers: quite a patch of this form, none of the ordinary green near it, though elsewhere on E.)	401 C E
174	C	405 C
176	C E S	407 C
178	C E	410 C E
182	E	418 C
198	C S	419 C
202	C	422 C E
213	E	429 C
214	C E S	430 C E
216	C E S	431 C E
217	C	439 C
219	E	445 C
244	C	446 Nordport, Aug 1887
248	C	449 C
249	C	462 C
264	C S	463 C E
265	O	468 C
269	C	493 C
270	C E S	501 C
271	C	505 C E
272	C E	506 E S
273	C	509 C E S
274	C	514 C E S
275	C E	522 C E
279	C	529 C E
280	C	534 E
283	C E	526 C E S
286	C E	540 C E S
		542 E

547	E		812	C.E.
548	E.	Persistent by old <del>gate</del> <sup>house, destroyed over 20 years ago.</sup>	820a	C.E.
553	C		822	C.E.
556	C.E.		824	C.E.
557	C.E.S.		825	C.E.
561	C.E.S.		827	E
567	C.E.S.		829	C. & bushing
576	C.E.		831	C.E.S.
579	C.E.		834	C.E.S.
580	C		836	C.E.
583	C		837	C.E.
584	C		846	C.E.
585	E		848	C.S.
591	C.E.S.		849	S
599	C		853	C
601	C.S.		854	C
601a	C		882	S
607	Trackport, Aug. 1887: not over 200 ft } elevation		895	C.E.
609	C.E.		896	S
613	C.E.		898	C.E.S.
623	C.E.		900	C.E.S.
627	C		905	C.S.
635	E		920	C.E.S.
637	E		942	C
641	S		943	C
644	C		945	C.E. down to sea level.
648	C.E.		952	C.E.S.
653	C		955	C.E.
654	C.E.		956	C.E.
663	E		958	C
664	C		959	C.E.
696	C.E.		966	C.E.
699	C		968	Trackport. Aug. 1887 (fruit)
701	S	along the shore	977	C.
706	C.E.S.		985	E
708	E		986	C
720	E		990	C
723	E		1001	C
725	E.S.		1008	C.E.S.
737	C.S.	also Cushing, islands in St. George's Bay.	1010	C.E.
740	E		1011	C. apparently rarer than 1010
761	C		1024	C
769	C		1025	C.E.
771	C.E.S.		1028	C
779	E.S.		1034	C
783	C		1033	C
786	C.S.		1046	C.E.
787	C.E.S.		1054	C.E.S.
790	C		1055	C.E.
793	C.E.S.		1058	E
798	C.E.S.		1067	C.E.
799	C.E.	common	1070	C
801	C.E.	chiefly in marshy places	1073	E
802	C.E.S.	" on rocks	1075	E
			1078	E.S.

1079	E. S.
1084	C
1090	C
1111	C E
1112	C E
1133	C
1135	E
1138	C E
1141	E
1143	E
1144	C E
1152	C
1156	E
1161	C
1164	Northport Aug. 1887
1177	C, also Cushing July 1888
1178	C
1179	Cushing July 1888
1186	E
1189	C E
1189b	C E
1191	C E
1198	E
1212	C E
1219b	C E
1232	C
1248	E
1261	E. S.
1274	C
1301	E
1309	C
1310	C
1320	C
1322	Northport Aug. 1887
1340	E
1342	C
1349	C E
1350	C
1359	C S
1366	C S
1368	C E
1370	C E
1371	C
1372	C S
1374a	C S
1376	C S
1383	C E S
1387	C S
1389	C S
1390	C S
1399	C
1402a	C
1403	C
1404	C

*Eupatorium purpureum*  
low grounds at C  
*Senecio viscosus*  
one plant near the shore  
& remote from any house. E.

*Glaux maritima*  
Rather common near  
high water mark. C

*Rhynchos raphaniticum*  
persistent near old houses at  
E. I saw large in  
plants, in blossom, ~~by~~  
at places where the houses  
had been destroyed & the  
garden deserted over 25  
years ago.